

## A Two Generation of Familial Adenomatous Polyposis

Lily Chandrawati\*, I Dewa Nyoman Wibawa\*\*

\*Department of Internal Medicine, Universitas Udayana/Sanglah General Hospital, Denpasar

\*\*Division of Gastroentero-hepatology, Department of Internal Medicine  
Universitas Udayana/Sanglah General Hospital, Denpasar

### Corresponding author:

I Dewa Nyoman Wibawa. Division of Gastroentero-hepatology, Department of Internal Medicine, Sanglah General Hospital, Jl. Diponegoro Dauh Puri Klod Denpasar Bali Indonesia. Phone/facsimile: +62-361-244177. Email: agusbobwibawa@yahoo.com

### ABSTRACT

Familial adenomatous polyposis (FAP) is a part of genetic polyposis syndrome which is caused by germline mutation in the adenomatous polyposis coli (APC) gene located in chromosome 5q21. The pathognomonic features is formation of hundreds to thousands of colorectal adenoma in late childhood and increase in size and number during adolescence. If left untreated, almost 100% patients will develop colorectal cancer by the age 50 years. We present a case of 26 year old male who complain of rectal bleeding, diarrhea, abdominal bloating, and has multiple polyps on colonoscopic finding. Two years ago, his father was diagnosed with polyposis coli and transverse colon adenocarcinoma. The patient was planned for preventive total colectomy. In conclusion, surgery remains the cornerstone treatment of FAP and surveillance program for early detection of cancer for all family member is very important to reduce colorectal cancer-related mortality.

**Keywords:** familial adenomatous polyposis (FAP), adenomatous polyposis coli (APC) gene, colorectal cancer.

### ABSTRAK

Familial adenomatous polyposis (FAP) merupakan salah satu bagian dari sindroma poliposis genetik yang disebabkan oleh mutasi gen adenomatous polyposis coli (APC) yang terletak pada kromosom 5q21. Gambaran klinis FAP yang khas adalah terbentuknya ratusan hingga ribuan adenoma pada usia akhir anak-anak dan adenoma ini cenderung bertambah besar dan banyak pada kelompok usia dewasa muda. Jika tidak mendapatkan penanganan, hampir 100% kasus berkembang menjadi karsinoma kolorektal pada usia 50 tahun. Berikut disajikan sebuah kasus, laki-laki, 26 tahun, dengan keluhan perdarahan rektum, diare, perut terasa kembung, dan dari gambaran kolonoskopi ditemukan adanya polip multipel di sepanjang kolon. Dua tahun yang lalu, ayah pasien didiagnosis polyposis coli dan adenokarsinoma kolon transversum. Pada pasien ini direncanakan untuk menjalani kolektomi total. Sebagai kesimpulan, pembedahan merupakan satu-satunya terapi FAP dan program surveilans untuk deteksi dini FAP pada seluruh anggota keluarga sangatlah penting untuk menurunkan mortalitas terkait karsinoma kolorektal.

**Kata kunci:** familial adenomatous polyposis (FAP), adenomatous polyposis coli (APC) gene, kanker kolorektal

## INTRODUCTION

Familial adenomatous polyposis (FAP) is a part of genetic polyposis syndrome which is caused by germline mutation in the adenomatous polyposis coli (APC) gene located in chromosome 5q21.<sup>1</sup> The incidence is estimated at one in every 8,000-10,000 live births and accounts for less than 1% of cases of colon cancer.<sup>2,3</sup> The frequency is constant throughout the world, with men and women equally affected.<sup>4</sup>

The clinical manifestation of FAP consists of intestinal and extraintestinal manifestation. Colonic adenoma represents the first intestinal manifestation, meanwhile the extraintestinal manifestations include congenital hypertrophy of retinal epithelium, epidermoid cyst, desmoid tumor, and pancreatic cancer.<sup>1,3</sup> Patients tend to develop hundreds to thousands of adenomatous polyps between the second and third decades of life.<sup>1</sup> If left untreated, almost 100% of patients will develop colorectal cancer by the age 50 years.<sup>3</sup>

Colorectal carcinoma (CRC) is the main cause of death among FAP patients, therefore early diagnosis and surveillance is imperative to reduce cancer-related mortality.<sup>3</sup> Surgery remains to be the definitive treatment of FAP, but the timing and type of surgery performed are still a controversy.<sup>1,5</sup>

## CASE ILLUSTRATION


A 26-year-old Balinese man presented with rectal bleeding, diarrhea, and abdominal bloating since 1 year. Two months ago he underwent rectal polypectomy in another private hospital. His father had similar complaints and was diagnosed with polyposis coli and adenocarcinoma of the transverse colon in 2016. On digital rectal examination, no palpable mass was found. His blood investigations were within normal limits. On colonoscopy, there were multiple polyps less than 1 cm in the entire colon from rectum to caecum, which is very suggestive of polyposis syndrome (Figure 1). Histopathological findings revealed adenomatous polyp with mild to moderate dysplasia (Figure 2). The patient was investigated for FAP-associated lesions and no other significant findings were noted. He was planned for preventive total colectomy with ileorectal anastomosis and his sister had been advised to do a colonoscopy screening.

## DISCUSSION

FAP, also known as adenomatous polyposis coli and familial polyposis coli, is the most common genetic polyposis syndrome. It comprises less than 1% of

colorectal cancer and is typically caused by an autosomal dominant germline mutation in the APC gene located on chromosome 5q21. Approximately up to one-third of cases appear to be caused by de novo mutation, without any clinical or genetic evidence of FAP in family members. The incidence is estimated at one in every 8,000-10,000 live births, with men and women equally affected.<sup>3,4,6</sup>


The clinical features of FAP can be divided into intestinal and extraintestinal manifestations. Colonic adenoma represents the first gastrointestinal manifestation and usually develops in the second or third decade of life with an average age of 15.9 years.<sup>3,7</sup> Classic FAP is characterized by hundreds to thousands of colorectal adenomatous polyps, while attenuated FAP develops fewer polyps, usually less than 100 and tends to occur later in life. Over time, the adenomas increase in size and number, so does the risk of cancer. Benign extraintestinal manifestations include congenital hypertrophy of retinal epithelium, epidermoid cyst, desmoid tumor, supernumerary teeth, osteoma, and adrenal adenoma. Other cancer risks are central nervous system, pancreatic, thyroid, and liver cancer.<sup>3,8</sup>




**Figure 1. Colonoscopic image showing multiple polyps in entire colon**

Herein two-generation cases of FAP are presented in which a son and his father had clinical presentations of FAP. Like most cases of FAP, these two cases showed an inherited pattern, which affects father and son in the family. Both of these cases had intestinal manifestations only without any extraintestinal signs. Diagnosis of polyposis syndrome was based on clinical findings with greater than 20 adenomas on colonoscopy and positive family history.<sup>9,10</sup> Over time, the risk of cancer is increasing and if left untreated, almost 100% of cases will develop colorectal cancer by the age of 50 years.<sup>3</sup> This is in line with his father's case who had colorectal adenocarcinoma at 50 (Figure 3). When patients are clinically diagnosed with FAP, they should undergo genetic testing in order to counsel their first-degree relatives regarding screening and treatment.<sup>9</sup> Unfortunately, the genetic testing mutation in this

case cannot be done because of limited facility in our hospital. Delay of diagnosis in our case maybe due to his father's long asymptomatic period. FAP has been a rare diagnosis in Indonesia and probably this above case was the first case documented in Indonesia, especially in Bali. FAP diagnosis is still become a challenge in our setting because of no national guidelines and suveillance has been established yet.



**Figure 2.** Histopathology revealed tubulovillous adenoma with mild to moderate dysplasia: 100x (2a) and 400x (2b) power magnification views of a hematoxylin-eosin stain.



**Figure 3.** Father's colonoscopy image showed multiple polyps (3a) and adenocarcinoma colon (3b)

Surgery remains the mainstay treatment of FAP. The primary goals of FAP treatment is cancer prevention and maximizing quality of life.<sup>11</sup> Multiple factors need to be considered when determining the proper operation including appropriate timing of surgery, extended area to be removed and the type of surgery to be performed. General agreement must always consider the individual characteristics of each patient.<sup>10</sup> Indication for early surgery include polyps > 10 mm diameter, polyps with high grade dysplasia, marked increase in polyp

number, and the present of symptoms. Surgical options are colectomy with ileorectal anastomosis (IRA), proctocolectomy with ileal pouch anal anastomosis (IPAA), and proctocolectomy with ileostomy.<sup>3,12,13</sup> Our case was planned for total colectomy with IRA. This approach was considered because of the low rectal polyp burden, has lower complication rate and better quality of life compared to IPAA.

CRC is the main cause of death among patients with FAP and surveillance leads to a reduction in CRC-related mortality. Colonoscopy screening should be performed in individual with clinical or genetic diagnosis of FAP. In those where no mutation can be found, all at risk relatives must undergo endoscopic screening. Colonoscopy should be begin at puberty or whenever the suggestive symptoms appeared. Colon screening guidelines for children with classic FAP is every 1-2 year sigmoidoscopy beginning at 10 to 12 years of age. For those who initially screening at older age should have colonoscopy for the first examination.<sup>3,14</sup> His sister was also advised for colonoscopy screening.

Prophylaxis colectomy remains the definitive therapy of FAP. Surveillance program for early detection of cancer for all at risk relatives is very important to reduce colorectal cancer-related mortality. Data concerning FAP in our country is scarce, compromising a better understanding about clinical aspects and treatment outcome. We hope in the future, the national registry could ameliorate this problem and improve medical care of FAP patients.

## REFERENCES

1. Feitosa MR, Fernandes de Oliveira THG, Kondo BRV, Gomes de Lira H, Abissamra AA, Para RS, et al. The epidemiological and clinical features of familial adenomatous polyposis in ribeiro preto. *J Coloproctol* 2013;33:126-30.
2. Plawski A, Banasiewicz T, Borun P, Kubaszewski L, Krokowicz P, Skrzypczak-Zielinska M, et al. Familial adenomatous polyposis of the colon. *Hered cancer Clin Pract* 2013;11:1-9.
3. Hagen CE, Setia N, Lauwers GY. Familial adenomatous polyposis: a review of gastrointestinal manifestation. *Diagnostic Histopathology* 2015;21:152-60.
4. Syngal S, Brand RE, Church JM, Giardiello FM, Hampel HL, Burt RW. ACG clinical guideline: genetic testing and management of hereditary gastrointestinal cancer syndrome. *Am J Gastroenterol* 2015;110:223-62.
5. Tudyka VN, Clark SK. Surgical treatment in familial adenomatous polyposis. *Ann Gastroenterol* 2012;25:201-6.
6. Dalavi SB, Vedpalsingh TH, Bankar SS, Ahmed MHS, Bhosale DN. Familial adenomatous polyposis (FAP) – a case study and review of literature. *J Clin Diagn Res* 2015;9:1-6.
7. Itzkowitz SH, Potack J. Colonic polyps and polyposis syndrome. In: Feldman M, Fiedman LS, Brandt LJ, eds.

Gastrointestinal and liver disease. 10<sup>th</sup> ed. Philadelphia: Elsevier Saunders 2016.p.2213-49.

8. Leoz ML, Carballal S, Moreira L, Ocana T, Balguer F. The genetic basis of familial adenomatous polyposis and its implications for clinical practice and risk management. *App Clin Genet* 2015;8:95-107.
9. Shussman N, Wexner SD. Colorectal polyps and polyposis syndrome. *Gastroenterology Report* 2014;2:1-15.
10. Herzig D, Hardimann K, Weiser M, Yu N, Paquette I, Feingold DL, et al. The american society of colon and rectal surgeons clinical practice guidelines for the management of inherititd polyposis syndromes. *Dis Colon Rectum* 2017;60:881-93.
11. Warriar SK, Kalady MF. Familial adenomatous polyposis: challenges and pitfalls of surgical treatment. *Clin Colon Rectal Surg* 2012;25:83-9.
12. Murgic J, Kirac I, Soldic Z, Tomas D, Zovak M, Bolanca A, et al. Familial adenomatous polyposis in three generation of a single family: a case study. *Case Rep Oncol* 2014;7:349-56.
13. Campos FG. Surgical treatment of familia adenomatous polyposis: dilemmas and current recommendations. *World J Gastroenterol* 2014;20:16620-9.
14. Mahajan NA, Mulay SS. Familial adenomatous polyposis of colon-a case report with review literature. *Int J Health Sci Res* 2015;5:388-92.